## 化学療法実施計画書(投与日程)

<u>レジメン名=胃癌TS-1+DOC</u> (対象疾病=

 《初回開始日》
 年 月 日

 今回[ ]クール目

身 長<u>=\_\_\_\_</u> 体 重=<u>\_\_\_</u> cm kg 体表面積= m<sup>2</sup> Ccr =

担当医

	薬剤	1日投与量	投与法	役与法 【投与時間 【1クール投与スケジュール(日目)																											
		$(/m^2 \text{ or } / \text{kg})$			1				5 6						12	13	14	15	16	17 1	8 19	9 20	0 21	22	23	24	25 2	6 27	28	29	30
	タキソテール(DOC:ドセタキセル)	40mg/ <b>m</b> ²	点静	60以上																		_	1								
_2	TS-1	80mg/ <b>m</b> ³*	経口		$\left[ ight]$	<b>〕</b> ]	<b>Ţ</b> ]	Ţ Į Į	ĮΊ	[[↓	<u>[</u> [1	<u> </u>	<u> </u>	]]	] ]	[↓[	$[\ ar{lackslash}\ ]$					]_	]_:								
l							_	_	.↓.	_  .	_ _	_	_ _		┨	ļ.,				_ _	_ _	_	┨	l			L .	_			
	体表面積	<u></u> <u>-</u> -				_	_	_	. ļ.	_  .	- L	_	_ _	<b>-</b>	┨	<b> </b>			_	- -	_ _	_ _	<b>.</b>	<b>.</b> .	 			_			
	★TS-1 1.25㎡未満	40mg×2回/日					-	_ \ _	<b>.</b>	_	-  -	- - :	- -	<b>-</b>	<b> </b>	4-4			- -	_ _	- -	_	4	<b> </b>				-			
	1.25~1.5㎡未満						- 🕹	- + -	. <b>.</b> .	-	- -	- - :	- -		┨	4 - 4	:	-	- -	- -	- -	_	┨	<b>.</b> .				-			
***	1.5㎡以上  5-HT₃拮抗薬	60mg×2回/日 1A	点静						+	+	-				-						+										_
					I-¦-∣		- +	- + -	-	-	- -	- - :	- -	-	┥	╅╾┪		-		- -	- -	- -	<b>-</b>	<b> </b>		-		-			
技遊	デキサメタゾン	3.3 <b>∼</b> 6.6mg	思朗		-⁴-		- +	- + -	-	-	- -	- - :	- -		┥- :		:	-		- -	- -	- -			+	-		-			
*							- †	- + -	- † -	- -	- -	- - :	- -		┨- :	1-1		-		- -	- -	-	<del> </del> -	<b>-</b> -	-	-		-			
投						Ī	Ť		Ť	Ť	<del>-  </del>	+	+	<del>†</del>	<del>†      </del>						Ť	<del>                                     </del>	+	Ī				i			ヿ
投与日程						- 1	- †	- † -	† †	- -	-  -	- -	- -		<b>1</b> - '	1-1	1		- -	- -	- -	-	1-	1	- 1		· -  - ·	-			- 1
Ιń						- 1	- †	- † -	† †	- -	-  -	- -	- -		1-	1-1			- -	- -	1-	1-	1-	1	-			-		- 1	-1
副									Ι.														<u> </u>	I							コ
作							_	_	.↓.	_  .	_ _	_	_ _		<b>.</b>	ļ.,				_ _	_ _		<u>.</u>	<b>.</b> .	↓			_			
用																															
		21日	クール回数																												
	休薬期間																—														
	☆予想される		白血球減少、好中球減少、口内炎、貧血、食欲不振 G-CSF投与、減量。																												
	重篤副作用																														
	 ☆副作用対策																														
	ス田リトのグラス		G-CSF投与、減重。 Cr>2.0、WBC<2,000																												
	☆実施中止条件	生化学項目																													
		副作用	, –- ,	· · - ·		-																									
		その他																													