

### 化学療法実施計画書（投与日程）

レジメン名＝膀胱癌・胆道癌TS-1+GEM

(対象疾病＝ )

《初回開始日》 年 月 日  
 今回〔 〕クール目

身長＝ \_\_\_\_\_ cm  
 体重＝ \_\_\_\_\_ kg  
 体表面積＝ \_\_\_\_\_ m<sup>2</sup>  
 Ccr = \_\_\_\_\_

担当医	

薬剤 (商品名, 一般名)	1日投与量 (/m <sup>2</sup> or /kg)	投与法 点静・静注	投与時間 (min)	1クール投与スケジュール(日目)																																		
				1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30					
1 ゲムシタビン(GEM)	1000mg/m <sup>2</sup>	点静	30	↓																																		
2 TS-1	80mg/m <sup>2</sup> *	経口		↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓																						
体表面積																																						
★TS-1 1.25m <sup>2</sup> 未満	40mg×2回/日																																					
1.25～1.5m <sup>2</sup> 未満	50mg×2回/日																																					
1.5m <sup>2</sup> 以上	60mg×2回/日																																					
前投薬	5-HT <sub>3</sub> 拮抗薬 デキサメタゾン	1A 点静 点静		↓																																		
投与日程																																						
副作用																																						
1クール期間	28日	クール回数																																				
休薬期間																																						
☆予想される 重篤副作用		骨髄抑制、食欲不振、下痢、口内炎など																																				
☆副作用対策		減量投与																																				
☆実施中止条件	生化学項目 副作用 その他	WBC<2000、Hb<8、Plt<70000、T.B>2.0、GOT>190、GPT>220、Cre>1.65																																				