

化学療法実施計画書（投与日程）

レジメン名=胆道癌TS-1+GEM(外科)

※外科患者限定レジメン

(対象疾病=)

身長= _____ cm

体重= _____ kg

体表面積= _____ m²

Ccr = _____

担当医	

《初回開始日》 _____ 年 _____ 月 _____ 日

今回(_____)クール目

薬剤 (商品名, 一般名)	1日投与量 (/m ² or /kg)	投与法 点静・静注	投与時間 (min)	1クール投与スケジュール(日目)																																	
				1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30				
1 ジェムザール(GEM)	1000mg/m ²	点静	30	↓																																	
2 TS-1	60mg/m ²	経口		↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓			
3																																					
4																																					
前投薬	デカドロン 5-HT3拮抗剤	10mg/body 1A	点静 点静	↓																																	
投与日程																																					
副作用																																					
1クール期間	28日	クール回数		*DLF=投与量規定因子																																	
休薬期間																																					
☆予想される重篤副作用		白血球減少、貧血(主に、ジェムザール) 食欲低下、眠気、口内炎、色素沈着など(主に、TS-1)																																			
☆副作用対策		*DLF=血液毒性 副腎皮質ホルモン、5-HT3拮抗剤の投与																																			
☆実施中止条件	生化学項目 副作用 その他	WBC<2000 (Seg<1000)															WBC<Grade2 (2000~2900) 血小板<Grade2 (50~74×103)																				
		重篤な血液毒性などの副作用															貧血 Grade2																				
		*JCOG副作用判定基準 Grade2																																			